



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

While Gauss refers to two earlier synthetic constructions of Paucker,¹ as the only previous ones which he knew had been publicly discussed, he remarks that that of Erchinger is “different and carried through more in the spirit of pure geometry.”

Stäckel reviews the discussions of the division of a circle into n equal parts, by means of imaginary quantities and the solution of the equation $x^n - 1 = 0$, by Cotes,² De Moivre,³ Euler,⁴ and Vandermonde,⁵ and the development of their ideas by Gauss in connection with the problem of constructing a regular inscribed polygon.

PROBLEMS FOR SOLUTION.

[N.B. The editorial work of this department would be greatly facilitated if, on sending in problems, the proposers would also enclose their solutions—*when they have them*. If a problem proposed is not original the proposer is requested *invariably* to state the fact and to give an exact reference to the source.]

2843. Proposed by E. H. MOORE, University of Chicago.

Show that the maximum of the absolute value of

$$2(a + ib)(x + iy) + i(a + ib)(z + iw) + i(c + id)(x + iy),$$

where $i = \sqrt{-1}$ and a, b, c, d, x, y, z, w are real numbers for which

$$a^2 + b^2 + c^2 + d^2 = x^2 + y^2 + z^2 + w^2 = 1,$$

is $1 + \sqrt{2}$. Study the locus of point-pairs $P = (a, b, c, d)$, $Q = (x, y, z, w)$ of the unit-sphere in real four-space for which this absolute value assumes its maximum value.

2844. Proposed by J. L. RILEY, Stephenville, Texas.

Decompose into simple fractions the number $\frac{60923880351}{1271888726}$ (Gauss, *Disq. Arith., Werke*, vol. 1, pp. 386–387).

2845. Proposed by E. L. POST, Princeton University.

Prove that if y_x is a solution of the functional equation

$$y_x = \frac{y_{x+1}^2}{x} + y_{x+1}$$

for positive integral values of x with $y_x > 0$, then

$$\lim_{x \rightarrow \infty} y_x \log x = 1.$$

2846.

Find the entire volume within the surface $x^{1/2} + y^{1/2} + z^{1/2} = a^{1/2}$. (W. A. Granville, *Elements of Differential and Integral Calculus*, revised ed., 1911, p. 420.)

This equation, rationalized, is the equation of Steiner's quartic surface, every tangent plane to which cuts it in two conics. (Cf. Salmon-Rogers, *Analytic Geometry of Three Dimensions*, 5th ed., vol. 2, 1915, pp. 171, 201, 207, 213f. Also C. M. Jessop, *Quartic Surfaces* 1916, Chapter 7.)

¹ (1) “Geometrische Verzeichnung des regelmässigen 17-Ecks und 257-Ecks in d. Kreis,” *Jahres-verhandl. d. kurländischen Gesellschaft für Litteratur und Kunst*, Mitau, Band 2, 1822. (2) *Die ebene Geometrie der geraden Linie und des Kreises*. Königsberg, 1823, p. 187. Paucker is also the author of: (3) *De divisione geometrica peripheriae circuli in XVII partes æquales*, Königsberg, 1817.

² R. Cotes, *Harmonia mensurarum, sive analysis et synthesis per rationum et angulorum mensuras promota*. Cambridge, 1722.

³ A. De Moivre, *Miscellanea analytica*, London, 1730.

⁴ L. Euler, *Introductio in analysin*, Lausanne, 1748, especially t. 1, cap 8: De quantitativibus transcendentibus ex circulo ortis.

⁵ C. A. Vandermonde, “Remarques sur les problèmes de situation,” *Histoire de l'Acad.*, année 1771, Paris 1774; *Mémoires*, p. 566. “Sur la résolution des équations,” p. 365.